

An application of Kleene's fixed point theorem to dynamic programming

Takashi Kamihigashi,* Kevin Reffett[‡] and Masayuki Yao[§]

We show that the least fixed point of the Bellman operator in a certain set can be computed by value iteration whether or not the fixed point is the value function. As an application, we give a simpler proof of one of Kamihigashi's main results.

Key words dynamic programming, Bellman equation, value function, fixed point

JEL classification C61

Accepted 9 July 2014

1 Introduction

Dynamic programming is one of the most important tools in studying dynamic economic models. Recently, under a minimal set of conditions, Kamihigashi (2014a) showed that the value function of a stationary dynamic programming problem can be computed by value iteration.

In this note we show that the least fixed point of the value iteration algorithm, or the Bellman operator, in a certain set can be computed whether or not the fixed point is the value function. Although it is insufficient for computing the value function itself, this result is significant in that it separates the issue of convergence, or computability, from the existence and uniqueness of a fixed point of the Bellman operator.

We prove the above result by applying what is known as “Kleene's fixed point theorem,” which has rarely been used directly in economics.¹ We present this fixed point theorem in the next section. In Section 3 we introduce some definitions and notation. In Section 4 we prove our main result. In Section 5 we illustrate the usefulness of our main result by showing one of the main results of Kamihigashi (2014a) with a simpler proof.

2 Kleene's fixed point theorem

In this section we present “Kleene's fixed point theorem” (e.g., Baranga 1991). We begin with some mathematical terminology. Let (P, \preceq) be a partially ordered set; i.e., \preceq is a reflexive, antisymmetric,

* Research Institute for Economics and Business Administration, Kobe University, Japan. Email: tkamihig@rieb.kobe-u.ac.jp

‡ Department of Economics, Arizona State University, USA.

§ Department of Economics, Keio University, Japan.

Financial support from the Japan Society for the Promotion of Science (KAKENHI 15H05729) and the Keio University Doctorate Student Grant-in-Aid Program 2013 is gratefully acknowledged. We would like to thank Hiroyuki Ozaki, Takako Fujiwara-Greve, Toru Hokari, Shinsuke Nakamura, Mikio Nakayama, and Chaowen Yu for helpful comments and suggestions.

¹ One exception is Vassilakis (1992).

and transitive binary relation on P . An *upper bound* of a set $Q \subset P$ is an element $p \in P$ satisfying $q \preceq p$ for all $q \in Q$. The *least element* of $Q \subset P$ is an element $p \in Q$ satisfying $p \preceq q$ for all $q \in Q$; note that the least element, if it exists, is uniquely defined since \preceq is antisymmetric.² The *supremum* of $Q \subset P$, denoted as $\sup Q$, is the least upper bound of Q .

A sequence $\{p_n\}_{n \in \mathbb{N}}$ is called *increasing* if $p_n \preceq p_{n+1}$ for all $n \in \mathbb{N}$. We say that (P, \preceq) is ω -*complete* if every increasing sequence in P has a supremum in P . A function $f : P \rightarrow P$ is called ω -*continuous* if for every increasing sequence $\{p_n\}$ in P having a supremum, we have $f(\sup_{n \in \mathbb{N}} p_n) = \sup_{n \in \mathbb{N}} f(p_n)$. An ω -continuous function f is necessarily increasing in the sense that $f(p) \preceq f(q)$ whenever $p \preceq q$.

Theorem 1 *Let (P, \preceq) be an ω -complete partially ordered set. Let $f : P \rightarrow P$ be ω -continuous. Suppose that there exists $\underline{p} \in P$ such that $\underline{p} \preceq f(\underline{p})$. Then $\underline{p}^* \equiv \sup_{n \in \mathbb{N}} f^n(\underline{p})$ is the least fixed point of f in $\{p \in P : \underline{p} \preceq p\}$.*

PROOF: See Stoltenberg-Hansen, Lindström, and Griffor (1994, p. 24). □

Baranga (1991) calls the above result “Kleene’s fixed point theorem,” while Sangiorgi (2009, p.35) states that “it is indeed unclear who should be credited for the theorem.”³ According to Jachymski, Gajek, and Pokarowski (2000, p. 249), Theorem 1 is equivalent to the Tarski–Kantorovitch fixed point theorem (e.g., Granas and Dugundji 2003, Theorem 1.2, p. 26).⁴

3 Dynamic programming

Our setup is identical to that of Kamihigashi (2014a). Here we briefly introduce the definitions and notation necessary for presenting our results.

Let X be a set, and Γ be a non-empty-valued correspondence from X to X . Let D be the graph of Γ :

$$D = \{(x, y) \in X \times X : y \in \Gamma(x)\}. \tag{1}$$

Let $u : D \rightarrow [-\infty, \infty)$. Let Π and $\Pi(x_0)$ denote the set of feasible paths and the set of feasible paths from x_0 , respectively:

$$\Pi = \{\{x_t\}_{t=0}^\infty \in X^\infty : \forall t \in \mathbb{Z}_+, x_{t+1} \in \Gamma(x_t)\}, \tag{2}$$

$$\Pi(x_0) = \{\{x_t\}_{t=1}^\infty \in X^\infty : \{x_t\}_{t=0}^\infty \in \Pi\}, \quad x_0 \in X. \tag{3}$$

Let $\beta \geq 0$. The value function $v^* : X \rightarrow \overline{\mathbb{R}}$ is defined by

$$v^*(x_0) = \sup_{\{x_t\}_{t=1}^\infty \in \Pi(x_0)} \mathop{\text{L}}\limits_{T \uparrow \infty} \sum_{t=0}^T \beta^t u(x_t, x_{t+1}), \quad x_0 \in X, \tag{4}$$

² That is, if $p \preceq q$ and $q \preceq p$, then $p = q$.

³ To be precise, Sangiorgi (2009) refers to a special case of Theorem 1.

⁴ The Tarski–Kantorovitch fixed point theorem has recently been used quite extensively in economics; see Mirman, Morand, and Reffett (2008), Van Zandt (2010), Balbus, Reffett, and Wozny (2013, 2014), and McGovern, Morand, and Reffett (2013), among others.

where $L \in \{\underline{\lim}, \overline{\lim}\}$ with $\underline{\lim} = \lim \inf$ and $\overline{\lim} = \lim \sup$. We define

$$\Pi^0 = \left\{ \{x_t\}_{t=0}^\infty \in \Pi : \underset{T \uparrow \infty}{L} \sum_{t=0}^T \beta^t u(x_t, x_{t+1}) > -\infty \right\}, \tag{5}$$

$$\Pi^0(x_0) = \{ \{x_t\}_{t=1}^\infty \in \Pi(x_0) : \{x_t\}_{t=0}^\infty \in \Pi^0 \}, \quad x_0 \in X. \tag{6}$$

Following the convention that $\sup \emptyset = -\infty$, we see that

$$\forall x_0 \in X, \quad v^*(x_0) = \sup_{\{x_t\}_{t=1}^\infty \in \Pi^0(x_0)} \underset{T \uparrow \infty}{L} \sum_{t=0}^T \beta^t u(x_t, x_{t+1}). \tag{7}$$

Let V be the set of functions from X to $[-\infty, \infty)$. The Bellman operator B on V is defined by

$$(Bv)(x) = \sup_{y \in \Gamma(x)} \{u(x, y) + \beta v(y)\}, \quad x \in X, v \in V. \tag{8}$$

A fixed point of B is a function $v \in V$ such that $Bv = v$.

The partial order \leq on V is defined in the usual way:

$$v \leq w \iff \forall x \in X, v(x) \leq w(x). \tag{9}$$

Given $v, w \in V$ with $v \leq w$, we define the order interval $[v, w]$ by

$$[v, w] = \{f \in V : v \leq f \leq w\}. \tag{10}$$

4 The main result

We are ready to prove the main result of this note as a consequence of Kleene's fixed point theorem.

Theorem 2 *Suppose that there exist $\underline{v}, \overline{v} \in V$ such that*

$$\underline{v} \leq \overline{v}, \tag{11}$$

$$B\underline{v} \geq \underline{v}, \tag{12}$$

$$B\overline{v} \leq \overline{v}. \tag{13}$$

Then the following conclusions hold:

- (a) $\underline{v}^* \equiv \sup_{n \in \mathbb{N}} (B^n \underline{v})$ is the least fixed point of B in $[\underline{v}, \overline{v}]$.
- (b) The increasing sequence $\{B^n \underline{v}\}_{n \in \mathbb{N}}$ converges to \underline{v}^* pointwise.

PROOF: By (11)–(13), B maps $[\underline{v}, \overline{v}]$ into itself. We apply Kleene's fixed point theorem to $B : [\underline{v}, \overline{v}] \rightarrow [\underline{v}, \overline{v}]$. For this purpose, it suffices to verify that $[\underline{v}, \overline{v}]$ is ω -complete and that B is ω -continuous on $[\underline{v}, \overline{v}]$.

To see that $[\underline{v}, \bar{v}]$ is ω -complete, let $\{v_n\}_{n \in \mathbb{N}}$ be an increasing sequence in $[\underline{v}, \bar{v}]$. Then the pointwise supremum of $\{v_n\}$ is equal to $\sup_{n \in \mathbb{N}} v_n$, the least upper bound of $\{v_n\}$. Since $\underline{v} \leq v_n \leq \bar{v}$ for all $n \in \mathbb{N}$, we have $\sup_{n \in \mathbb{N}} v_n \in [\underline{v}, \bar{v}]$. It follows that $[\underline{v}, \bar{v}]$ is ω -complete.

To see that B is ω -continuous on $[\underline{v}, \bar{v}]$, let $\{v_n\}_{n \in \mathbb{N}}$ be an increasing sequence in $[\underline{v}, \bar{v}]$ again. Let $x \in X$. We have

$$[B(\sup_{n \in \mathbb{N}} v_n)](x) = \sup_{y \in \Gamma(x)} \{u(x, y) + \beta(\sup_{n \in \mathbb{N}} v_n)(y)\} \tag{14}$$

$$= \sup_{y \in \Gamma(x)} \sup_{n \in \mathbb{N}} \{u(x, y) + \beta v_n(y)\} \tag{15}$$

$$= \sup_{n \in \mathbb{N}} \sup_{y \in \Gamma(x)} \{u(x, y) + \beta v_n(y)\} \tag{16}$$

$$= [\sup_{n \in \mathbb{N}} (Bv_n)](x), \tag{17}$$

where (15) holds since $u(x, y)$ is independent of n , and (16) follows by interchanging the two suprema using Kamihigashi (2008, Lemma 1). Since x is arbitrary, it follows that $B \sup_{n \in \mathbb{N}} v_n = \sup_{n \in \mathbb{N}} Bv_n$. That is, B is ω -continuous on $[\underline{v}, \bar{v}]$.

Now by Kleene's fixed point theorem, conclusion (a) follows. To see conclusion (b), note from (12) that $\underline{v} \leq B\underline{v} \leq B^2\underline{v} \leq \dots$. Thus $\{B^n \underline{v}\}_{n \in \mathbb{N}}$ is increasing, and for each $x \in X$, we have $(B^n \underline{v})(x) \uparrow \underline{v}^*(x)$ as $n \uparrow \infty$ by the definition of \underline{v}^* ; i.e., $\{B^n \underline{v}\}$ converges to \underline{v}^* pointwise. \square

Theorem 2 allows one to compute the least fixed point of B in $[\underline{v}, \bar{v}]$ by iterating B from \underline{v} . Of course, this fixed point may not be the value function v^* without additional conditions, but it is exactly this point that makes Theorem 2 significant: convergence to the least fixed point is valid whether or not it is the value function.

5 Applications

In this section, we illustrate the usefulness of Theorem 2 by showing how to use it to prove one of the main results of Kamihigashi (2014a), which replaces \underline{v}^* with the value function v^* in the conclusions of Theorem 2.

The next lemma slightly generalizes an argument used in the proof of Theorem 4.3 in Stokey and Lucas (1989), providing a sufficient condition for any fixed point v of B with $\underline{v} \leq v$ to satisfy $v^* \leq v$.⁵

Lemma 1 *Let $\underline{v} \in V$ be such that*

$$\forall \{x_t\}_{t=0}^\infty \in \Pi^0, \quad \liminf_{t \uparrow \infty} \beta^t \underline{v}(x_t) \geq 0. \tag{18}$$

Let $v \in V$ be a fixed point of B with $\underline{v} \leq v$. Then $v^ \leq v$.*

PROOF: See the Appendix. \square

The following result is immediate from Theorem 2 and Lemma 1.

⁵ Although the lemma is not explicitly shown in Kamihigashi (2014a), it is recognized and mentioned in Kamihigashi (2014a,b).

Lemma 2 (Kamihigashi 2014a, Lemma 6.3). Let $\underline{v}, \bar{v} \in V$ satisfy (11)–(13) and (18). Then $v^* \leq \underline{v}^*$, where \underline{v}^* is defined in Theorem 2.

This result is proved in Kamihigashi (2014a) using the additional result that finite iterations of B correspond to finite-horizon approximations of the infinite-horizon problem (4) (Kamihigashi 2014a, Lemma 6.2). Our proof of Lemma 2 shows that this approximation result is unnecessary once Theorem 2 and Lemma 1 are available. Let us now use Theorem 2 and Lemma 2 to prove one of the main results of Kamihigashi (2014a).⁶

Theorem 3 (Kamihigashi 2014a, Theorem 2.2). Suppose that $v^* \in V$. Suppose further that there exists $\underline{v} \in V$ satisfying (11), (18), and

$$\underline{v} \leq v^*. \quad (19)$$

Then the following conclusions hold:

- (a) v^* is the least fixed point of B in $[\underline{v}, \bar{v}]$.
- (b) The increasing sequence $\{B^n \underline{v}\}_{n=1}^\infty$ converges to v^* pointwise.

PROOF: Since v^* is a fixed point of B (see Kamihigashi 2014a, Lemma 2.1), (11)–(13) hold with $\bar{v} = v^*$. Define \underline{v}^* as in Theorem 2. Then $\underline{v}^* \leq \bar{v} = v^*$. The reverse inequality holds by Lemma 2; thus $\underline{v}^* = v^*$. Now both conclusions follow from Theorem 2. \square

As another application of Theorem 2, it can be used along with Lemma 6.1 in Kamihigashi (2014a) to show Theorem 2.1 in Kamihigashi (2014a).

Appendix

Proof of Lemma 1

We show that $v^*(x_0) \leq v(x_0)$ for any $x_0 \in X$. Let $x_0 \in X$. If $\Pi^0(x_0) = \emptyset$, then $v^*(x_0) = -\infty \leq v(x_0)$. Suppose that $\Pi^0(x_0) \neq \emptyset$. Let $\{x_t\}_{t=1}^\infty \in \Pi^0(x_0)$. We have

$$\begin{aligned} v(x_0) &\geq u(x_0, x_1) + \beta v(x_1) \\ &\geq u(x_0, x_1) + \beta u(x_1, x_2) + \beta^2 v(x_2) \\ &\quad \vdots \\ &\geq \sum_{t=0}^{T-1} \beta^t u(x_t, x_{t+1}) + \beta^T v(x_T) \\ &\geq \sum_{t=0}^{T-1} \beta^t u(x_t, x_{t+1}) + \beta^T \underline{v}(x_T). \end{aligned}$$

⁶ In Kamihigashi (2014a, Theorem 2.2), v^* is shown to be the least fixed point of B in a larger set. The same set can be used here with an additional argument.

Applying $L_{T \uparrow \infty}$ to the right-hand side, we have

$$v(x_0) \geq \underset{T \uparrow \infty}{L} \left\{ \sum_{t=0}^{T-1} \beta^t u(x_t, x_{t+1}) + \beta^T \underline{v}(x_T) \right\} \tag{A1}$$

$$\geq \underset{T \uparrow \infty}{L} \sum_{t=0}^{T-1} \beta^t u(x_t, x_{t+1}) + \overline{\lim}_{T \uparrow \infty} \beta^T \underline{v}(x_T) \tag{A2}$$

$$\geq \underset{T \uparrow \infty}{L} \sum_{t=0}^{T-1} \beta^t u(x_t, x_{t+1}), \tag{A3}$$

where (A2) follows from the properties of $\overline{\lim}$ and $\underline{\lim}$,⁷ and (A3) holds by (18). Since (A1)–(A3) hold for any $\{x_t\}_{t=1}^\infty \in \Pi^0(x_0)$, applying $\sup_{\{x_t\}_{t=1}^\infty \in \Pi^0(x_0)}$ to (A3) and recalling (7), we obtain $v(x_0) \geq v^*(x_0)$. \square

References

Baranga, A. (1991), “The contraction principle as a particular case of Kleene’s fixed point theorem,” *Discrete Mathematics* **98**, 75–9.

Balbus, and L., K. Reffett, and L. Wozny (2013), “A qualitative theory of large games with strategic complementarities,” mimeo, Arizona State University.

Balbus, L., K. Reffett, and L. Wozny (2014), “A constructive study of Markov equilibrium in stochastic games with strategic complementarities,” *Journal of Economic Theory* **150**, 815–40.

Granas, A., and J. Dugundji (2003), *Fixed Point Theory*, New York: Springer-Verlag.

Jachymski, J., L. Gajek, and P. Pokarowski (2000), “The Tarski–Kantorovitch principle and the theory of iterated function systems,” *Bulletin of the Australian Mathematical Society* **61**, 247–61.

Kamihigashi, T. (2008), “On the principle of optimality for nonstationary deterministic dynamic programming,” *International Journal of Economic Theory* **4**, 519–25.

Kamihigashi, T. (2014a), “Elementary results on solutions to the Bellman equation of dynamic programming: Existence, uniqueness, and convergence,” *Economic Theory* **56**, 251–73.

Kamihigashi, T. (2014b), “An order-theoretic approach to dynamic programming: An exposition,” *Economic Theory Bulletin* **2**, 13–21.

McGovern, J., O. F. Morand, and K. Reffett (2013), “Computing minimal state space recursive equilibrium in OLG models with stochastic production,” *Economic Theory* **54**, 623–74.

Michel, P. (1990), “Some clarifications on the transversality condition,” *Econometrica* **58**, 705–23.

Mirman, L., O. F. Morand, and K. Reffett (2008), “A qualitative approach to Markovian equilibrium in infinite horizon economies with capital,” *Journal of Economic Theory* **139**, 75–98.

Sangiorgi, D. (2009), “On the origins of bisimulation and coinduction,” *ACM Transactions on Programming Language and Systems* **31**, article 15.

Stokey, N., and R. E. Lucas, Jr. (1989), *Recursive Methods in Economic Dynamics*, Cambridge, MA: Harvard University Press.

Stoltenberg-Hansen, V., I. Lindström, and E. R. Griffor (1994), *Mathematical Theory of Domains*, Cambridge: Cambridge University Press.

Van Zandt, T. (2010), “Interim Bayesian Nash equilibrium on universal type space for supermodular games,” *Journal of Economic Theory* **145**, 249–63.

Vassilakis, S. (1992), “Some economic applications of Scott domains,” *Mathematical Social Sciences* **24**, 173–208.

⁷ We have $\underline{\lim}(a_t + b_t) \geq \underline{\lim}a_t + \underline{\lim}b_t$ and $\overline{\lim}(a_t + b_t) \geq \overline{\lim}a_t + \overline{\lim}b_t$ for any sequences $\{a_t\}$ and $\{b_t\}$ in $[-\infty, \infty)$ whenever both sides are well defined (e.g., Michel, 1990, p. 706).